

**EFFECTING OF SUCROSE
CONCENTRATIONS AND INOCULUM
DENSITY ON ADVENTITIOUS ROOT
GROWTH IN CELL SUSPENSION CULTURE
OF *Panax vietnamensis* AND INITIALLY
GROWTH IN A BIOREACTOR**

**Nguyen Van Ket*, Truong Thi Lan Anh
and
Nguyen Hoang Uyen Dung**

*Department of Agroforestry,
Dalat University - Vietnam
e-mail: ketnv@dlu.edu.vn*

Abstract

It has been demonstrated that the difference of initial sucrose concentrations (0, 3, 5, 7, 9 %) effected on adventitious root growth in cell suspension culture of *Panax vietnamensis*. The maximum adventitious root growth was obtained in a concentration of 5% sucrose. The density of initial explants also effected to the course of growth. The concentration of 30 mM.2% inoculum density was optimal, and further increase of inoculum density led to a specific decrease of the rate root growth. Adventitious root of *Panax vietnamensis* developed well in 5 - liter bioreactors containing 2 liters of SH medium (working medium) supplemented with 5 mg l⁻¹ IBA, and 0,02 mg/l BA. The growth rate of root was 5.6 folds after 45 days in this condition.

*Corresponding author

Key words: Vietnamese ginseng, sucrose concentrations, density, bioreactor

1. Introduction

This is a specific variety of Vietnamese ginseng, called Sâm Ngoc Linh (*Panax vietnamensis* Ha et Grushv). It was found in high mountains of Vietnam central at the height from 1200 - 2100 metres above sea level. The southern central area such as Ngoc Lum Heo and Ngoc Am (Quang Nam) and highlands such as Ngoc Linh (Dakto), Tra Mi are key distributions of *Panax*. It is considered a highly valuable medicine used for the treatments of various diseases, exciting nervous systems, stimulating memories and immune systems, enhancing vitality and hormone, supporting anti-oxidant and anti-depression, regulating heart blood circulation, reducing blood cholesterol content and body inflammation. As a result, Ngoc Linh ginseng has been exploring dramatically and it becomes scarce in nature and highly extinctive if no actively protective treatments have been desired.

In fact, in recent years, various researches on mass propagation in nature conditions have been conducted. However, the pharmaceutical market demand of Ngoc Linh ginseng has been so big that powder and extracts from the mountain collection has not been enough due to its extreme slow growth. That is the result why more researches have been recently attracted to speed up Ngoc Linh ginseng production. In addition to cell tissue culture in vitro, cell suspension culture have been made. The methods were also applied to other varieties of ginseng such as *Panax ginseng* (Yu, 2000 v Thanh et al., 2005), *Panax notoginseng* (Zhang et al., 1995). In reality, researches on adventitious root growth of Sâm Ngoc Linh have been very few and need being continued in order to find out a potential alternative to a huge commercial markets worldwide of ginseng.

2. Materials and Methods

1. Materials Adventitious roots of Ngoc Linh ginseng from callus at 0.5 - 1cm.
2. Culture condition: The samples were cultivated in darkness at room temperature 25°C.
3. Methods: - Generating and quick propagation of adventitious roots SH medium (Schenk and Hildebrandt) supplemented 5mg/l IBA and 8% agar (data report is not preferred) has been used as induction medium for generating and quick propagation of Ngoc Linh adventitious roots from callus.
 - The effect of sucrose concentrations to culture mediumAventitious root lines at different density rates 2, 4, 6, 8 % were maintained SH medium (Schenk and Hildebrandt) supplemented 5mg/l IBA with various sucrose concentrations. The root growth rate was checked after 45 days of cultivation.
 - Cultivation in bioreactors Adventitious roots were cut into small sections

of 1-1.5cm and then cultivated in 5 - liter bioreactors containing 2 liters of SH medium (working medium) supplemented 5 mg l-1 IBA, and 5% sucrose concentration. Root lines were maintained in darkness at room temperature 25 20C. Fresh weight and dry weight were done after 45 days of cultivation.

- The effect of cultivated density Adventitious roots were cultivated in 100 ml conical flasks with SH medium (working medium) supplemented 5 mg l-1 IBA, and 50 g sucrose.

3. Results and Discussion

3.1 Results

After 45 days of culture, the final fresh weight and dry weight were increased considerably with the increase of initial sucrose concentrations from 0 to 5%. Higher sucrose concentrations (7-9%) seemed to repress adventitious root growth (Table 1). In fact, culture condition at 5% sucrose concentration is optimized for the adventitious root growth of Ngoc Linh ginseng (3378 mg fresh weight and 344 mg dry weight).

Table 1: The effect of sucrose concentrations to the root growth of Ngoc Linh ginseng (*Panax vietnamensis*). The data collected after 45 days of culture in 100 ml conical flasks with a 20 ml working volume of SH medium.

Sucrose concentrations %	Fresh weight (mg)	Dry weight (mg)
0	1034 ^d	49 ^c
3	2514 ^c	278 ^b
5	3378 ^a	344 ^a
7	2859 ^b	336 ^a
9	2309 ^c	326 ^a
CV%	5,13	4,68
LCD _{0,05}	225,90	22,74

The effects of cultured density Different culture densities increasing from 2, 4, 6, 8 % were investigated. After 45 days, the results indicated that culture at 2% resulted in highest growth rate; and other increasings of cultured densities were observed in relatively decreasing growth rate. Truly, the optimum culture density for adventitious root growth of Ngoc Linh ginseng was at 2%, 5.3 times higher than initial samples (Table 2).

Table 2: The effect of different culture densities to the growth of adventitious roots of Ngoc Linh ginseng (*Panax vietnamensis*). The data was collected

after 45 days of culture in 100 ml conical flasks with a working volume of 20 ml SH medium.

Densities%(w/v)	Fresh weight(mg)	Dry weight(mg)	Growth rate
2	1732 ^d	190 ^d	4.3 ^a
4	3010 ^c	298 ^c	3.8 ^b
6	3740 ^b	349 ^b	3.1 ^c
8	4198 ^a	395 ^a	2.6 ^d

The culture in bioreactors Adventitious roots started increasing in length and inducing callus after 10 days of culture. New adventitious roots changed in color, relatively light yellow, after 21 days. And after 45days, fresh weight of 45 g was obtained, making an increase of 5.6 times higher than initial samples.

3.2 Discussion

3.2.1 Effect of sucrose concentrations Sucrose has been used as optimim carbon source to plant cell and tissue cultures (Giaquinta, 1980; Strickland et al. 1987). Sucrose supplies carbons to the process of plant growth as well as energy to cell life. Similarly, sucrose also provides essential carbons to the formation of bi-products from metabolism. In a sufficient medium with essential nutritions, an increase of sucrose concentraton potentially affected to an increase of biomass and the ratio of the metabolism progress for bi-products (Rokem v Goldberg, 1985).

Within this experiment, increasing of innital sucrose concentrations from 0 to 5% was studied and shown that after 45 days of culture fresh weight and dry weight were increased accordingly (Table 1).

It gives the same result as culture of *Panax ginseng* (Choi et al., 1994a, 1994b; Akalezi, 1999). On the other hand, further increase of sucrose concentrations from 7 to 9% led to a heavily negative effect to root growth rate. It can be said that penetration potential of cell membrance made a certain negative effect to root growth and to secondary metabolite cell production (Narayan, 2002). When studying *Panax notoginseng*, Zhang (1995), it was also said that higher sucrose cncentrations (above 5%) inhibited rooth growth significantly.

3.2.2 The effect of different culture densities The initial culture density showed a considerable effect to the developing of cell suspension. A lower initial culture density lessened cell division. It is likely that the substance from cells of cell suspension was developing that stimulated cell growth. In fact, the initial cell density plays a specific role, namely 'an adapter', promoted the growth of cell suspension (Henshaw et al., 1966; Torees, 1989). Consequently, culture


density is tremendously important in the process of sub-cultures of plant cells and similarly affected to the progress of bi-products (Rokem v Goldberg, 1985; Liang et al., 1991). On the contrary, a relatively high initial culture density will lessen the growth of cell suspension due to competition between nutrients and oxygen source from culture medium (Henshaw et al, 1966; Torees, 1989). Based on the experiment, it can be concluded that optimal density to root growth is made by 2%; and it is apparent that the growth steadily decreases at higher culture densities. The result is similar to the report of the above named authors.

3.2.3 Culture of *Panax vietnamensis* adventitious roots in bioreactors

Traditional plant cell culture methods for mass production of ginseng still encounters various constraints because of uncapable adjustment of physical and chemical factors in the cultured flasks and of expensively manual labour cost. Therefore, application of culture in bioreactors is getting more beneficial in in-vitro culture, is much easier to respectively adjust to optimal conditions for the growth of adventitious roots (increasing of gas exchange and of output of metabolic production process), is able to produce an adequate yield throughout the year and makes a lower production cost (Levin et al., 1988; Preil et al., 1988, 1991; Sharma, 1992; Christie et al., 1995; Honda et al., 2001; Paek et al., 2001). In principle, culture of ginseng in bioreactor chains, essential nutrients can be regularly provided and supplemented to adventitious roots especially when the culture medium becomes exhausted and unables to supply adequate nutrition to the samples. For this reason, culture in bioreactors is beneficial to better growth of samples (Yao v Zhong, 1999). In addition, culture in bioreactors, due to a continual gas supplement source, adventitious roots are able to get relatively higher oxygen than that from a traditional cell culture. As a result, root systems grew up vigorously. Supplement of oxygen source affected to the growth and secondary metabolic synthesis in various plant cultures including *Catharanthus roseus* (Leckie et al., 1991) v *Perilla frutescens* (Zhong et al., 1993).

Cell cultures in liquid medium *Thalictrum minus* to produce berberine, cells that synthesized berberine obtained oxygen content two times higher than that did not (Kobayashi et al., 1991). Cultures in liquid medium and continual supplement of gas provide an adequate oxygen content for the growth and secondary metabolic synthesis.

The experiment showed that after 10 days of culture, adventitious roots started developing in size and inducing callus, and that after 21 days of culture, new adventitious roots became light yellow. After 45 days of culture, fresh weight of adventitious roots were obtained by 45 g and growth rate is 5.6 times higher than the samples. After that, such adventitious got older and seemed to change in colour, getting darker. Then, subculture needed to be done respectively.



Picture 1: The process of biomass production of roots of Ngoc Linh ginseng.

a) Fresh mountain ginseng roots collected from Ngoc Linh Mountain; b) Adventitious Ngoc Linh ginseng roots developed from callus; c) Roots cultured in 100 ml conical flasks; d) Roots cultured in bioreactors.

4. Conclusion

Adventitious roots of Ngoc Linh P. Vietnamensis cultured in liquid medium after 45 days, our results are as follows:

- The optimal concentration of sucrose for adventitious root growth was by 5%;
- The optimal density for adventitious root growth was by 2% and other concentrated densities inhibited adventitious root growth.
- After 45 days of culture, adventitious roots of Ngoc Linh ginseng cultured in 5-liter bioreactors with a working volume of 2 liter SH medium supplemented 5mg/l IBA developed 5.6 times higher than innital samples.

References

- [1] Akalezi C.O., Liu S., Liu Q.S., Yu J.T., Zhong J.J. (1999). Combined effects of initial sucrose concentration and inoculum size on cell growth and ginseng saponin production by suspension cultures of *Panax ginseng*. *Pro. Biochem.* 34: 639-642.
- [2] Choi I.C.Y., Ahn I.O., Park J.C. (1994a). Production of ginseng saponin in tissue culture of ginseng (*Panax ginseng* C.A. Mayer). *Russ. Jour. Plant Physio.* 140:784-788.
- [3] Choi K.T., Lee C.H., Ahn I.O., Lee J.H., Park J.C. (1994b). Characteristics of the growth and ginsenosides in the suspension culture cells of Korean ginseng (*Panax ginseng* C. A. Mayer). In Bailey, Whitehead C., Proctor J.T.A , Kyle J.T., eds, *Proc Intl Ginseng Conf.*, Vancouver B.C., Canada, pp 259-268.
- [4] Christie J.A., Kozai T., Takayama S. (1995). Automation in plant tissue culture: general introduction and overview. In *Automation and environmental control in plant tissue culture*. Christie
- [5] J.A., Kozai T., Smith M.L., eds. Kluwer Academic Publishers, Dordrecht Do Huy Bich et al. (2003). *Plants and animals using as medicines in Vietnam*, Vol II. Science, Natural Sciences and Technology Publisher, Ha Noi.
- [6] Giaquinta R.T. (1980). Translocation of sucrose and oligosaccharides. In: *Press Jour. (ed) Biochemistry of Plants*, Vol. 3: 271-320. Academic Press, Inc.
- [7] Honda H., Liu C., Kobayashi Y. (2001). Large scale plant propagation. In: *Advances in Biochemical Engineering/Biotechnology*. Vol., 72. Scheper T, ed. Springer - Verlag Berlin Heidelberg
- [8] Kobayashi Y., Fukui H., Tabata M. (1991). Enhancement of berberine production by spermidine in *Thalictrum minus* cell suspension cultures. *Plant Cell Rep* 9:496.
- [9] Leckie F., Scragg A.H., Cliffe K.C. (1991). An investigation into the role of initial KLa on the growth and alkaloid accumulation by cultures of *Catharanthus roseus*. *Biotechnol. Bioeng.*, 37: 364 - 370
- [10] Levin R., Gaba V., Tal B., Hirsh S., De Nola D., Vasil K. (1988). Automated plant tissue culture for mass propagation. *Biotechnology* 6: 1035 -1040
- [11] Liang S.Z., Zhong J.J., Yoshida T. (1991). Review of plant cell culture technology for producing useful products (part 1). *Ind. Microbiol*, 21 (3): 27 - 31 (in China)
- [12] Narayan M. S., Venkataraman, L. V. (2002). Effect of Sugar and Nitrogen on the Production of Anthocyanin in Cultured Carrot (*Daucus carota*) cells. *Journal of Food Science*, vol. 67: 84 - 86. Nguyen Tien Ban et al. (2007). *Vietnamese Red Book*. Part II. Botany. Science, Natural Sciences and Technology Publisher, Ha Noi.
- [13] Nguyen Thi Lieu., Nguyen Trung Thanh., Nguyen Van Ket (2011). The adventitious root induce of Ngoc Linh ginseng (*Panax vietnamensis*, Ha et Grushv.) in vitro cultures. *VNU Journal of Science, natural Sciences and Technology* 27 (2011) 30-36.
- [14] Nguyen Trung Thanh, Nguyen Van Ket, Paek Kee Yoeup. Effecting of medium composition on biomass and ginsenoside production in cell suspension culture of *Panax vietnamensis* Ha et Grushv. *VNU Journal of Science, natural Sciences and Technology* 23 (2007) 269-274
- [15] Paek K.Y., Hahn E. J., Son S. H. (2001). Application of Bioreactors for Large-Scale Micropropagation Systems of Plants. *In Vitro Cell. Dev. Biol-Plant* 37: 149-157.
- [16] Preil W. (1991). Application of bioreactors in plant propagation. In: Debergh PC and Zimmerman RH (eds.), *Micropropagation*, kluwer Academic Publisher, Netherlands pp 425 - 445 Preil W., Florek P., Wix U., Beck A. (1988). Towards mass propagation by use of bioreactors. *Acta Hort* 226: 99 - 105
- [17] Rokem J.S., Goldberg I. (1985). Secondary metabolites from plant cell suspension cultures: Methods for yield improvement. In: *Advances in Biotechnological Processes* 4 (Mizrahi A, Wezel AL, eds). Alan R. Liss, New York. 241 - 274

- [18] Sharma N.K. (1992). Automation in plant tissue culture: problems and prospects. *Curr. Sci.*, 7: 507 - 512
- [19] Strickland S.G., Nichol J.W., Call C.M. and Stuart D.A. (1987). Effect of carbohydrate source on alfalfa somatic embryogenesis. *Plant Sci.* 48: 113-121
- [20] Thanh N.T., Murthy H.N., Yu K.W., Hahn E.J., Paek K.Y. (2005). Methyl jasmonate elicitation enhanced synthesis of ginsenoside by cell suspension cultures of *Panax ginseng* in 5-l balloon type bubble bioreactors. *Applied Microbiology and Biotechnology*, 67, 197-201.
- [21] Yao H., Zhong J.J. (1999). Improved production of ginseng polysaccharide by adding conditioned medium to *Panax notoginseng* cell cultures. *Biotechnology Techniques* 13: 347-349.
- [22] Yasuhiro F. and Yasuhiro H. (1985). Effective Production of Shikonin by Cultures with an Increased Cell Population. *Agric. Biol. Chem.*, 49 (7): 2071 -2075. Yu K.W., Hahn E.J. & Paek K.Y. (2000) Production of adventitious ginseng roots using bioreactors. *Kor. J. Plant Tiss. Cult.* 27: 309-315
- [23] Zhang Y.H., Zhong J.J., Yu J.T. (1995). Effect of osmotic pressure on cell growth and production of ginseng saponin and polysaccharide in suspension cultures of *Panax notoginseng*. *Biotechnology Letters*, vol. 17 (12): 1347-1350.
- [24] Zhong J.J., Yoshida M., Fujiyama K., Yoshida T. (1993). Enhancement of anthocyanin production by *Perilla frutescens* cells in a stirred bioreactor with internal light irradiation. *J. Ferment. Bioeng.*, 75: 299 - 303